

**Note to readers with disabilities:** *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact [ehp508@niehs.nih.gov](mailto:ehp508@niehs.nih.gov). Our staff will work with you to assess and meet your accessibility needs within 3 working days.

## **Supplemental Material**

# **Airborne Fine Particles and Risk of Hospital Admissions for Understudied Populations: Effects by Urbanicity and Short-Term Cumulative Exposures in 708 U.S. Counties**

Mercedes A. Bravo, Keita Ebisu, Francesca Dominici, Yun Wang, Roger D. Peng, and Michelle L. Bell

### **Table of Contents**

**Table S1.** Characteristics of PM<sub>2.5</sub> pollution data.

**Table S2.** Summary statistics of model evaluation for 24-h average PM<sub>2.5</sub> county level exposure estimates

**Figure S1.** Availability of monitor data by county for the study area output (data from 2000 U.S. Census; map created using ArcGIS).

**Figure S2.** County-specific correlation between exposure estimates derived from observed data and CMAQds simulated PM<sub>2.5</sub> concentrations for the 418 counties with monitoring data and populations  $\geq 50,000$ , 2002-2006 (data from 2000 U.S. Census; map created using ArcGIS). This map shows correlations between county level monitor-derived daily exposure estimates for daily county level CMAQds-derived exposure estimates. Only counties with monitoring data and populations  $\geq 50,000$  are included in the comparison

**Figure S3.** Comparison of county-specific maximum likelihood health effect estimates obtained from monitor-derived exposure estimates (x-axis) versus CMAQds\_subset-derived exposure estimates ( $\hat{\beta}^c$  coefficients relating  $PM_{2.5}$  concentration to hospitalization rates in county-specific regression model): (a) cardiovascular; (b) respiratory (n=418 counties) Point size is inversely proportional to the magnitude of the standard error associated with each monitor-derived county-specific maximum likelihood health effect estimate, such that a smaller point size indicates greater uncertainty associated with that county-specific estimate. Note that these values represent coefficients that have not been scaled.

**Figure S4.** Percent increase in hospital admissions associated with a  $10\mu g/m^3$  increase in  $PM_{2.5}$  concentration, estimated using monitoring data (gray) and downscaler output (black), only for counties with monitoring data (CMAQds\_subset), by level of urbanicity (lag 0) Vertical lines represent 95% posterior intervals. Urbanicity is measured as percent of county population residing in nonurban areas

## Tables

**Table S1. Characteristics of PM<sub>2.5</sub> pollution data/output**

	<b>Monitoring data</b>	<b>CMAQds output</b>
<b>Data description</b>	Federal Reference Method ambient air quality monitors	Daily predictions of pollutant concentrations at Census Tracts centroids from combination of ambient monitoring data and CMAQ v4.6 output
<b>Spatial form of concentration observation/estimate</b>	Point	Point
<b>Spatial resolution of original dataset</b>	Variable	Variable
<b>Temporal resolution</b>	Variable, ~1 observation/3 days	Daily, every day
<b>Method(s) used to estimate county level concentration</b>	Monitor(s) within given county averaged	Population weighted Census Tracts to estimate county level conc.
<b>Spatial coverage of exposure estimates</b>	~418 counties	~2,818 counties


**Table S2. Summary statistics of model evaluation for 24-h average PM<sub>2.5</sub> county level exposure estimates<sup>a,b</sup>**

<b>EvaluationMetric</b>	<b>Value</b>
Mean daily county level concentration	
CMAQds	12.28 µg/m <sup>3</sup>
CMAQds_subset	12.60 µg/m <sup>3</sup>
Observed (monitor-derived)	12.48 µg/m <sup>3</sup>
Normalized mean bias (NMB) (%)	0.95%
Normalized mean error (NME) (%)	9.75%
Mean correlation (standard deviation)	0.97 (0.032)


<sup>a</sup> Formulas and further description of metrics of model performance are presented in Zhang et al. 2006.

<sup>b</sup> The mean correlation refers to the mean correlation between monitor-derived and CMAQds-derived exposure estimates within a county (and not correlations across all counties and days).


## Figures




**Figure S1. Availability of monitor data by county for the study area (data from 2000 U.S. Census; map created using ArcGIS).**



**Figure S2. County-specific correlation between exposure estimates derived from observed data and CMAQs simulated PM<sub>2.5</sub> concentrations for the 418 counties with monitoring data and populations  $\geq 50,000$ , 2002-2006 (data from 2000 U.S. Census; map drawn using ArcGIS).** This map shows correlations between county level monitor-derived daily exposure estimates for daily county level CMAQs-derived exposure estimates. Only counties with monitoring data and populations  $\geq 50,000$  are included in the comparison



**Figure S3. Comparison of county-specific maximum likelihood health effect estimates obtained from monitor-derived exposure estimates (x-axis) versus CMAQs\_subset-derived exposure estimates ( $\hat{\beta}^c$  coefficients relating  $\text{PM}_{2.5}$  concentration to hospitalization rates in county-specific regression model): (a) cardiovascular; (b) respiratory (n=418 counties).** Point size is inversely proportional to the magnitude of the standard error associated with each monitor-derived county-specific maximum likelihood health effect estimate, such that a smaller point size indicates greater uncertainty associated with that county-specific estimate. Note that these values represent coefficients that have not been scaled.



**Figure S4. Percent increase in hospital admissions associated with a 10 $\mu$ g/m<sup>3</sup> increase in PM<sub>2.5</sub> concentration, estimated using monitoring data (gray) and downscaler output (black), only for counties with monitoring data (CMAQds\_subset), by level of urbanicity (lag 0). Vertical lines represent 95% posterior intervals. Urbanicity is measured as percent of county population residing in nonurban areas.**



## References

Zhang Y, Liu P, Pun B, Seigneur C. 2006. A comprehensive performance evaluation of MM5-CMAQ for the Summer 1999 Southern Oxidants Study episode - Part I: Evaluation protocols, databases, and meteorological predictions. *Atmos Environ* 40: 4825-4838.